## § 522.1372

(iii) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[68 FR 68724, Dec. 10, 2003, as amended at 69 FR 69523, Nov. 30, 2004]

## §522.1372 Mepivacaine.

- (a) Specifications. Each milliliter (mL) of solution contains 20 milligrams mepivacaine hydrochloride.
- (b) Sponsor. See No. 000009 ir  $\S510.600(c)$  of this chapter.
- (c) Conditions of use in horses—(1) Amount. For nerve block, 3 to 5 mL; for epidural anesthesia, 5 to 20 mL; for intra-articular anesthesia, 10 to 15 mL; for infiltration, as required; for anesthesia of the laryngeal mucosa prior to ventriculectomy, by topical spray, 25 to 40 mL, by infiltration, 20 to 50 mL.
- (2) Indications for use. For use as a local anesthetic for infiltration, nerve block, intra-articular and epidural anesthesia, and topical and/or infiltration anesthesia of the laryngeal mucosa prior to ventriculectomy.
- (3) Limitations. Not for use in horses intended for human consumption. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

## § 522.1380 Methocarbamol injection.

[71 FR 39547, July 13, 2006]

- (a) Specifications. The product is a sterile, pyrogen-free solution, each milliliter containing 100 milligrams of methocarbamol, 0.5 milliliter of polyethylene glycol 300, and water for injection q.s. Its pH is 3.5 to 6.0.
- (b) *Sponsor*. See No. 000856 in §510.600(c) of this chapter.
- (c) Conditions of use—(1) Amount—(i) Dogs and cats. 20 milligrams per pound of body weight for moderate conditions, 25 to 100 milligrams per pound of body weight for severe conditions (tetanus and strychnine poisoning), total cumulative dose not to exceed 150 milligrams per pound of body weight.
- (ii) Horses. 2 to 10 milligrams per pound of body weight for moderate conditions, 10 to 25 milligrams per pound of body weight for severe conditions (tetanus), additional amounts may be needed to relieve residual effects and to prevent recurrence of symptoms.
- (2) Indications for use. As an adjunct for treating acute inflammatory and

traumatic conditions of the skeletal muscles and to reduce muscular spasms.

(3) Limitations. For intravenous use only. For dogs, administer rapidly half the estimated dose, pause until the animal starts to relax, then continue administration to effect. For horses, administer rapidly to effect. Not for horses intended for food use. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[45 FR 79758, Dec. 2, 1980, as amended at 46 FR 18964, Mar. 27, 1981; 67 FR 67521, Nov. 6, 2002]

## § 522.1410 Methylprednisolone.

- (a) Specifications. Each milliliter of suspension contains 20 or 40 milligrams (mg) of methylprednisolone acetate.
- (b) *Sponsors*. See Nos. 000009 and 000010 in §510.600(c) of this chapter.
  - (c) [Reserved]
- (d) Conditions of use—(1) Dogs—(i) Amount. Administer 2 to 40 mg (up to 120 mg in extremely large breeds or dogs with severe involvement) by intramuscular injection or up to 20 mg by intrasynovial injection.
- (ii) Indications for use. For treatment of inflammation and related disorders; treatment of allergic and dermatologic disorders; and as supportive therapy to antibacterial treatment of severe infections.
- (iii) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- (2) Cats—(i) Amount. Administer 10 to 20 mg by intramuscular injection.
- (ii) Indications for use. For treatment of inflammation and related disorders; treatment of allergic and dermatologic disorders; and as supportive therapy to antibacterial treatment of severe infections.
- (iii) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- (3) *Horses*—(i) *Amount*. Administer 200 mg by intramuscular injection or 40 to 240 mg by intrasynovial injection.
- (ii) *Indications for use*. For treatment of inflammation and related disorders.